

## INFLUENCE OF MENTAL MOTOR IMAGERY ON THE EXECUTION OF A FINGER-TO-THUMB OPPOSITION TASK

A. GEMIGNANI<sup>1,2</sup>, M. DI STEFANO<sup>2</sup>, L. SEBASTIANI<sup>2</sup>, B. GHELARDUCCI<sup>2</sup>,  
M. JEANNEROD<sup>3</sup>, M. GUAZZELLI<sup>4</sup> AND R. MASSARELLI<sup>1,5</sup>

<sup>1</sup> INSERM U438, CHU de Grenoble, BP 217, 38043 Grenoble cedex 09, France; <sup>2</sup> Dipartimento di Fisiologia e Biochimica, Università di Pisa, 56127 Pisa, Italy; <sup>3</sup> Institut des Sciences Cognitives du CNRS, 67 Bvd Pinel, 69675 Bron, France; <sup>4</sup> Dipartimento di Psichiatria, Neurobiologia, Farmacologia e Biotecnologia, Università di Pisa, 56100 Pisa, Italy; <sup>5</sup> CNRS ERS 2019, Bat. R. Dubois, Campus la Doua, 69622 Villeurbanne cedex, France

### INTRODUCTION

The condition in which a subject consciously images and simulates a motor performance has been widely exploited (for a review see 12). Functional magnetic resonance imaging studies (fMRI) (20) and electrophysiological data (15) indicate a wide activation of the sensorimotor cortex during simulation of the movements, and a cerebellar activation during the mental imagery of complex motor actions (18). Motor imagery has also been associated with an increase of the premotor and prefrontal activity suggesting that the mental representation of the movements plays an important role in both planning and preparation stages of the performance (11). Recent fMRI reports point out that the same cortical regions are activated by pure motor imagery and by simple execution of movements (21, 22). There is however evidence that the pattern of cortical activity elicited by pure mental simulation differs from that occurring when motor imagery precedes the actual performance and it may be further changed by the mode of movements execution (4). These findings are suggestive of task-dependent functional interactions, between different neural structures involved in motor output and visuomotor imagination.

Taking into account these observations one may ask whether the mental rehearsal of movements performed immediately prior of the motor outcome may influence the pattern of activity of motor areas, compared to the simple execution. The present fMRI study addresses this question by comparing regional distribution of cortical activity during unilateral hand movements and during the same movements preceded by their mental simulation.




Fig. 1. - Schematic drawing of the experimental paradigm.

On the top is represented the E condition, simple Execution of the motor task, consisting in a resting period (R), followed by the motor outcome (E).

On the bottom is represented the S+E condition, Execution preceded by mental Simulation of the motor task, consisting in a resting period (R) followed by the mental simulation of the task (S) and by the actual execution of the simulated movements (E).

adigm (22). The z-scores of temporal responses were also calculated. The functional maps included only clusters comprising at least three adjacent "activated" voxels, which exhibited a cross-correlation factor with the paradigm  $> 0.7$  and a z-score  $> 0.5$  (2). Isolated activated voxels were discarded.

A blood vascular mask, derived from the three-dimensional PC MR scan, was used to identify possible macrovascular responses within the slices. In each subject the vascular masks were superimposed on the corresponding activation maps to rule out signal increments due to blood flow increase in the cortical macrovascular system (24). Finally, the functional images were superimposed onto the corresponding high-resolution anatomical images (derived from the three-dimensional PC MR scan) to localize the cortical activations. The identification of activated cortical regions was performed by visual comparison of the functional maps with the anatomical images of the Talairach and Tournoux atlas (25).

In each subject we studied three distinct cortical motor areas: primary motor cortex corresponding to the anterior bank of central sulcus (M1, Brodmann's Area 4); pre-motor area that spans from the precentral sulcus to the dorsolateral frontal lobe and includes the anterior precentral gyrus (PMA, lateral portion of Brodmann's Area 6); supplementary motor area that extends from the paracentral sulcus to cover the anterior region of the medial caudal half of superior frontal gyrus (SMA, medial portion of Brodmann's Area 6).

#### Statistical analysis

The consistency of cortical activation across the different runs was verified by ANOVA, having as main factors the number of voxels of each Scan, the Subjects and the Conditions. The results showed that under both the E and the S+E conditions there was no difference among the three runs, therefore the statistical analyses were performed on the total number of activated voxels measured in each subject. The data were analyzed by three-factor ANOVA for repeated measures having as




Fig. 2. - Functional activity in the two hemispheres.

Histograms show the total number of correlated voxels in the contralateral and ipsilateral hemisphere in the E condition (white columns) and the S+E condition (grey columns).

\* S+E vs E condition  $p < .05$ .

## DISCUSSION

The present experiment compares the functional cortical activity elicited by the execution of a finger-to-thumb opposition task (E condition) with the activity occurring when mental simulation of the task precedes the actual performance (S+E condition). We have found that the spatial distribution of activated voxels is increased in the S+E condition with respect to the E condition. This suggests that motor areas are functionally more active when the movements are performed after their mental simulation, relative to the condition without mental rehearsal.

Previous studies have reported that the pure motor imagery (19,22) as well as the simple execution of unimanual motor task (22) induces functional activation of the premotor areas (PMA) of both hemispheres. Accordingly, one may expect that the enhancement of activity occurring during the S+E condition is present on the ipsilateral as on the contralateral PMA. Our findings, while confirming the bilateral activation of PMA during the E condition, show that mental simulation has a rather different effect on the two sides of the brain. Indeed, functional activation of the contralateral PMA is significantly increased during the S+E condition with respect to

limitation. Several reports indicate in fact that the activity of the left motor cortex during ipsilateral fingers movements is more consistent than the corresponding ipsilateral activation of the right cortex (6, 14). Given this hemispheric functional asymmetry, one may expect that the mental imagery does influence the activation of ipsilateral MI at a different extent in the two hemispheres.

Our findings are in agreement with those results suggesting that the level of neural excitability may be increased by the motor imagery. Bonnet *et al.* (3) found increased spinal reflexes during a mentally simulated isometric foot pressure and similar results were obtained by using transcranial magnetic stimulation (TMS) of motor cortex. TMS was used to trigger motor evoked potentials (MEPs) in arm muscles during simulated arm movements and MEPs were increased only in the muscles involved in the imagined movements. (5, 8, 13). A logical consequence of increased motor cortex excitability is that it should propagate down to the motoneuron level. This is still a controversial issue, however (12).

All together these results suggest that motor performance preceded by mental rehearsal is associated to selective increases of the cortical activity. Not all the motor areas that are functionally active in simple motor execution and pure motor imagery present the enhancement of activity during the S+E condition. Changes in the pattern of cortical activity might reflect the concurrent activation of regions involved in both the preparation and the execution stage of the motor outcome. The activity change may be dependent on different factors, such as the subject's familiarity with task, the type of motor imagery and the mode of movements execution.

In conclusion, the beneficial effects of "mental practice" on the physical performance (26, 28) may rely on the increment of cortical activity due to the close temporal association between motor rehearsal and actual performance. The effect may be accounted by a sort of neural recruiting that is made possible by overlapping of the cortical networks involved in motor output and in motor imagery (19). Finally, the fact that the motor pathways are globally activated during motor imagery represents a rationale for rehearsing effects observed in normal subjects during the motor learning (23, 27) and opens new possibilities for the rehabilitation of patients with motor impairments (9).

#### SUMMARY

The present fMRI study compares regional distribution of the cortical activity during the execution of unilateral hand movements (finger-to-thumb opposition) preceded or not by their motor simulation (S+E and E condition, respectively). The results show that, overall, the number and the spatial distribution of activated voxels are both increased in the S+E with respect to the E condition. The motor performance preceded by mental rehearsal is related to selective increase of the cortical activity. Among the motor areas that are found active during the simple motor execution a significant enhancement of functional activation during the S+E condition has been found only in the contralateral premotor area (PMA) and in the contra-and

16. MAAS, L.C., FREDERICK, B.D. AND RENSHAW, P.F. Decoupled automated rotational and translational registration for functional MRI time series data: the DART registration algorithm. *Magn. Reson. Med.*, **37**: 131-139, 1997.
17. OLDFIELD, R.C. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*, **9**: 97-113, 1971.
18. PARSONS, L.M., FOX, P.T., DOWNS, J.H., GLASS, T., HIRSCH, T.B., MARTIN, C.C., JERABEK, P.A. AND LANCASTER, J.L. Use of implicit motor imagery for visual shape discrimination as revealed by PET. *Nature*, **375**: 54-58, 1995.
19. PORRO, C.A., CETTOLO, V., FRANCESCATO, M.P. AND BARALDI, P. Ipsilateral involvement of primary motor cortex during motor imagery. *Eur. J. Neurosci.*, **12**: 3059-3063, 2000.
20. PORRO, C.A. AND CORAZZA, R. Functional magnetic resonance imaging as a tool for investigating human cortical motor function. *Arch. Ital. Biol.*, **137**: 101-114, 1999.
21. PORRO, C.A., FRANCESCATO, M.P., CETTOLO, V., DIAMOND, M.E., BARALDI, P., ZUANI, C., BAZZOCCHI, M. AND DI PRAMPERO, P.E. Primary motor and sensory cortex activation during motor performance and motor imagery: a functional magnetic resonance imaging study. *J. Neurosci.*, **16**: 7688-7698, 1996.
22. ROTH, M., DECETY, J., RAYBAUDI, M., MASSARELLI, R., DELON-MARTIN, C., SEGEBARTH, C., MORAND, S., GEMIGNANI, A., DECORPS, M. AND JEANNEROD, M. Possible involvement of primary motor cortex in mentally simulated movement: a functional magnetic resonance imaging study. *Neuroreport*, **7**: 1280-1284, 1996.
23. ROURE, R., COLLET, C., DESCHAUMES-MOLINARO, C., DITTMAR, A., RAQA, H., DELHOMME, G. AND VERNET-MAURY, E. Autonomic nervous system responses correlate with mental rehearsal in volleyball training. *Eur. J. Appl. Physiol.*, **78**: 99-108, 1998.
24. SEGEBARTH, C., BELLE, V., DELON, C., MASSARELLI, R., DECETY, J., LE BAS, J.F., DECORPS, M. AND BENABID, A.L. Functional MRI of the human brain: predominance of signals from extracerebral veins. *Neuroreport*, **5**: 813-816, 1994.
25. TALAIRACH, J. AND TOURNOUX, P. *Co-planar stereotaxic atlas of the human brain*. Thieme, New York, 1988.
26. VOGT, S. On relations between perceiving, imagining and performing in the learning of cyclical movement sequences. *Br. J. Psychol.*, **86**: 191-216, 1995.
27. YAGÜEZ, L., NAGEL, D., HOFFMAN, H., CANAVAN, A.G.M., WIST, E. AND HÖMBERG, V. A mental route to mental learning. Improving trajectorial kinematics through imagery training. *Behav. Brain Res.*, **90**: 95-106, 1998.
28. YUE, G. AND COLE, K.J. Strength increases from the motor program: comparison of training with maximal voluntary and imagined muscle contractions. *J. Neurophysiol.*, **67**: 1114-23, 1992.

## INFLUENCE OF MENTAL MOTOR IMAGERY ON THE EXECUTION OF A FINGER-TO-THUMB OPPOSITION TASK

A. GEMIGNANI<sup>1,2</sup>, M. DI STEFANO<sup>2</sup>, L. SEBASTIANI<sup>2</sup>, B. GHELARDUCCI<sup>2</sup>,  
M. JEANNEROD<sup>3</sup>, M. GUAZZELLI<sup>4</sup> AND R. MASSARELLI<sup>1,5</sup>

<sup>1</sup> INSERM U438, CHU de Grenoble, BP 217, 38043 Grenoble cedex 09, France; <sup>2</sup> Dipartimento di Fisiologia e Biochimica, Università di Pisa, 56127 Pisa, Italy; <sup>3</sup> Institut des Sciences Cognitives du CNRS, 67 Bvd Pinel, 69675 Bron, France; <sup>4</sup> Dipartimento di Psichiatria, Neurobiologia, Farmacologia e Biotecnologia, Università di Pisa, 56100 Pisa, Italy; <sup>5</sup> CNRS ERS 2019, Bat. R. Dubois, Campus la Doua, 69622 Villeurbanne cedex, France

### INTRODUCTION

The condition in which a subject consciously images and simulates a motor performance has been widely exploited (for a review see 12). Functional magnetic resonance imaging studies (fMRI) (20) and electrophysiological data (15) indicate a wide activation of the sensorimotor cortex during simulation of the movements, and a cerebellar activation during the mental imagery of complex motor actions (18). Motor imagery has also been associated with an increase of the premotor and prefrontal activity suggesting that the mental representation of the movements plays an important role in both planning and preparation stages of the performance (11). Recent fMRI reports point out that the same cortical regions are activated by pure motor imagery and by simple execution of movements (21, 22). There is however evidence that the pattern of cortical activity elicited by pure mental simulation differs from that occurring when motor imagery precedes the actual performance and it may be further changed by the mode of movements execution (4). These findings are suggestive of task-dependent functional interactions, between different neural structures involved in motor output and visuomotor imagination.

Taking into account these observations one may ask whether the mental rehearsal of movements performed immediately prior of the motor outcome may influence the pattern of activity of motor areas, compared to the simple execution. The present fMRI study addresses this question by comparing regional distribution of cortical activity during unilateral hand movements and during the same movements preceded by their mental simulation.




Fig. 1. - Schematic drawing of the experimental paradigm.

On the top is represented the E condition, simple Execution of the motor task, consisting in a resting period (R), followed by the motor outcome (E).

On the bottom is represented the S+E condition, Execution preceded by mental Simulation of the motor task, consisting in a resting period (R) followed by the mental simulation of the task (S) and by the actual execution of the simulated movements (E).

adigm (22). The z-scores of temporal responses were also calculated. The functional maps included only clusters comprising at least three adjacent "activated" voxels, which exhibited a cross-correlation factor with the paradigm  $> 0.7$  and a z-score  $> 0.5$  (2). Isolated activated voxels were discarded.

A blood vascular mask, derived from the three-dimensional PC MR scan, was used to identify possible macrovascular responses within the slices. In each subject the vascular masks were superimposed on the corresponding activation maps to rule out signal increments due to blood flow increase in the cortical macrovascular system (24). Finally, the functional images were superimposed onto the corresponding high-resolution anatomical images (derived from the three-dimensional PC MR scan) to localize the cortical activations. The identification of activated cortical regions was performed by visual comparison of the functional maps with the anatomical images of the Talairach and Tournoux atlas (25).

In each subject we studied three distinct cortical motor areas: primary motor cortex corresponding to the anterior bank of central sulcus (M1, Brodmann's Area 4); pre-motor area that spans from the precentral sulcus to the dorsolateral frontal lobe and includes the anterior precentral gyrus (PMA, lateral portion of Brodmann's Area 6); supplementary motor area that extends from the paracentral sulcus to cover the anterior region of the medial caudal half of superior frontal gyrus (SMA, medial portion of Brodmann's Area 6).

#### Statistical analysis

The consistency of cortical activation across the different runs was verified by ANOVA, having as main factors the number of voxels of each Scan, the Subjects and the Conditions. The results showed that under both the E and the S+E conditions there was no difference among the three runs, therefore the statistical analyses were performed on the total number of activated voxels measured in each subject. The data were analyzed by three-factor ANOVA for repeated measures having as

ipsilateral primary motor regions (M1). The activity increase may be accounted by a sort of neural recruiting that is made possible by the overlapping of cortical networks involved in both motor output and motor imagery. The beneficial effects of "mental practice" on the physical performance may rely to the close temporal association between motor rehearsal and actual performance.

*Acknowledgments.* – The helpful contribution of Dr. D. Montanaro in the cortical localization of functional activations is gratefully acknowledged.

#### REFERENCES

1. BACIU, M.V., RUBIN, C., DECORPS, M.A. AND SEGEBARTH, C.M. fMRI assessment of hemispheric language dominance using a simple inner speech paradigm. *NMR. Biomed.*, **12**: 293-298, 1999.
2. BANDETTINI, P.A., JESMANOWICZ, A., WONG, E. C. AND HYDE, J.S. Processing strategies for time-course data sets in functional MRI of the human brain. *Magn. Reson. Med.*, **30**, 161-173, 1993.
3. BONNET, M., DECETY, J., REQUIN, J. AND JEANNEROD, M. Mental simulation of an action modulates the excitability of spinal reflex pathways in man. *Cogn. Brain. Res.*, **5**: 221-228, 1997.
4. DEIBER, M.P., IBANEZ, V., HONDA, M., SADATO, N., RAMAN, R. AND HALLETT, M. Cerebral processes related to visuomotor imagery and generation of simple finger movements studied with positron emission tomography. *Neuroimage*, **7**: 73-85, 1998.
5. FADIGA, L., BUCCINO, G., CRAIGHERO, L., FOGASSI, L., GALLESE, V. AND PAVESI, G. Corticospinal excitability is specifically modulated by motor imagery. A magnetic stimulation study. *Neuropsychologia*, **37**: 147-158, 1999.
6. HAALAND, K.Y. AND HARRINGTON, D.L. Hemispheric asymmetry of movement. *Curr. Opin. Neurobiol.*, **6**: 796-800, 1996.
7. HALL, C.R. AND PONGRAC, J. *Movement imagery questionnaire*. Faculty of Physical Education. The University of Western Ontario, London, Canada, 1983.
8. HASHIMOTO, R. AND ROTHWELL, J.C. Dynamic changes in corticospinal excitability during motor imagery. *Exp. Brain. Res.*, **125**: 75-81, 1999.
9. JACKSON, P.L., LAFLEUR, M.F., MALOUIN, F., RICHARDS, C. AND DOYON, J. Potential role of mental practice using motor imagery in neurologic rehabilitation. *Arch. Phys. Med. Rehabil.*, **82**: 1133-1141, 2001.
10. JEANNEROD, M. Mental imagery in the motor context. *Neuropsychologia*, **33**: 1419-1432, 1995.
11. JEANNEROD, M. AND DECETY, J. Mental motor imagery: a window into the representational stages of action. *Curr. Opin. Neurobiol.*, **5**: 727-732, 1995.
12. JEANNEROD, M. AND FRAK, V. Mental imaging of motor activity in humans. *Curr. Opin. Neurobiol.*, **9**: 735-739, 1999.
13. KASAI, T., KAWAI, S., KAWANISHI, M. AND YAHAGI, S. Evidence for facilitation of motor evoked potentials (MEPs) induced by motor imagery. *Brain. Res.*, **744**: 147-150, 1997.
14. KIM, S.G., ASHE, J., HENDRICH, K., ELLERMANN, J.M., MERKLE, H., UGURBIL, K. AND GEORGOPOULOS, A.P. Functional magnetic resonance imaging of motor cortex: hemispheric asymmetry and handedness. *Science*, **261**: 615-617, 1993.
15. LANG, W., CHEYNE, D., HOLLINGER, P., GERSCHLAGER, W. AND LINDINGER, G. Electric and magnetic fields of the brain accompanying internal simulation of movement. *Brain. Res. Cogn. Brain. Res.*, **3**: 125-129, 1996.






Fig. 3. - Mean activity change.

Difference between the number of activated voxels during S+E condition relative to the E condition for each region of interest of both hemisphere.

For each subject and for each experimental condition the activated clusters were localized according to the x, y, and z coordinates of the Talairach and Tournoux Atlas. The range of coordinates is indicated for M1, PMA and SMA of both hemispheres. Contralateral hemisphere: M1 (+32/+40, -20, +45/+65); PMA (+30, +15, +45/+65); SMA (+8/+10, -1/-20, +45/+65). Ipsilateral hemisphere: M1 (-32/-40, -20, +45/+65); PMA (-30, +15, +45/+65); SMA (-8/-10, -1/-20, +45/+65).

\* S+E vs E condition =  $p < .05$

the simple motor execution whereas the activity of ipsilateral PMA tends to be reduced in most of the subjects.

During the S+E condition the activity of primary motor cortex (M1) is significantly enhanced on both sides of the brain. The rate of change – that is estimated in relation to the activity in the E condition – is highest for the ipsilateral M1. The functional activation of ipsilateral M1 is rather poor during the E condition and becomes more than 7 times greater when the motor outcome follows the mental simulation of the task. In contralateral M1 the increment of activity during the S+E condition is smaller, although the total number of activated voxels is much higher than in ipsilateral M1.

Finally, the activity of supplementary motor area (SMA) during the S+E condition is virtually unchanged in both hemispheres with respect to E condition.

Our experiment does not allow a comparison between ipsilateral activity in the two hemispheres. The task was performed by the right hand only, and then ipsilateral activation could be measured only in the right hemisphere. This represents a

